

Removal of Lead Ions from Polluted Water Using *Plantago major* L. and *Phaseolus vulgaris* L.

Akram A. Ali and Ali A. AL-Homaidan

Department of Botany and Microbiology, Faculty of Science,
King Saud University, P.O. Box: 2455, Riyadh 11451, Kingdom of Saudi Arabia.

Abstract: This study focuses on lead detoxification using roots, stems, leaves and whole plants of *Plantago major* L. and *Phaseolus vulgaris* L. (decoration kidney bean) as a ball hanging hydroponically in deionized water supplemented with concentrations of lead ion under different duration times and temperatures regimes. Roots of studied plants showed that the highest removal of lead than other parts, while using a mixture of whole plants were more efficient than using each one alone. The largest amounts of lead 85, 77 and 71 % were accumulated in roots of *P. major* + *P. vulgaris*, *P. major* alone and *P. vulgaris* alone, respectively. The highest rate of Pb ions uptake from the medium took place during the first 2-days of incubation and after 5-days of incubation, lead content in the medium decreased by half. This study suggested that roots of *P. major* + *P. vulgaris* could be used removal of pollutants from water contaminated with lead.

Keywords: Detoxification; lead ions; hyperaccumulative plants, biosorption

INTRODUCTION

Human activities lead to Increase concentrations of heavy metals like lead in soil, water, air and living organisms caused by human activities (Seregin and Ivanov 2001). Elevated levels of heavy metals in water are taken up by the root system, accumulate in different parts of plants, reduce their growth and impair plant metabolism (Allinson and Dzialo 1981). On the other hand, introduction of these metals to man and animals through drinking water will help in occurring many diseases to them. Lead is one of the most widely distributed heavy metals and is very toxic to plants. Its effect on plants has been studied by several research terms (Bazzaz *et al.*, 1974; Titov *et al.*, 1996), but most work has been done on isolated systems.

The hyperaccumulator plants include about 400 species (Morel *et al.*, 1997). They are well-known and usually show the ability to accumulate 1000 times more heavy metal (lead) than normal plants without visible changes in their appearance or yield (Baker, 1995). At the same time, the level of lead accumulation can exceed even several hundred times the level permissible for living organism. Thus, lead will be introduced into the food chain and biological circulation. Therefore, the use of non-edible plants is preferable (Kucharski *et al.*, 1999).

The strategies of lead tolerance in plants are poorly understood. The tolerant ecotypes survive in an environment contaminated with lead and adopt mechanisms of avoidance and inactivation. Inactivation of lead, as well as of other heavy metals, is connected with the detoxification mechanism (Maitani *et al.*, 1996). Very often *P. major* and *P. vulgaris* plants grow in soil containing a high Pb concentration. From the results of various studies, Sheoran and Singh (1993) and Grill *et al.* (1989) have proposed several sites in the electron transport chain that are influenced by heavy metal. In the whole plant, Pb can affect photosynthesis at the stomatal level, mesophyll cells, pigment content, and light and dark reactions. Maitani *et al.* (1996) have shown that it affects chlorophyll biosynthesis in barley seedlings. At a high Pb content in soil, photosynthesis can also be reduced due to both a lower carboxylase activity and the effects on metabolites of the carbon reduction cycle (Stiborova *et al.*, 1986, 1987). As a result of reduced photosynthesis, the effect of Pb is perceived as a decrease in plant growth and development. A direct effect of Pb on stomata was described by Carlson *et al.* (1995). Studies on the use of air dried balls from *P. major* and *P. vulgaris* are new. These plants widely distributed in

Corresponding Author: Akram A. Ali, Department of Botany, Faculty of Science, Zagazig University, Zagazig, Egypt.
E-mail: akram692000@yahoo.com

natural and cultivated habitats. Different strategies for lead in environmental stress conditions by one or more than one organs of them may occur. To clarify which parts of the plant control the stability to increased Pb concentrations in water, we have studied its influence on the concentration of lead in water. This study is divided into three stages: first stage to select the best concentration from lead ion treatments; second is to determine the best duration to plant samples to take up the biggest amount; the last to show the effect of different temperatures on Pb-ion absorption.

MATERIALS AND METHODS

Experimental and Treatments:

Seeds of *Plantago major* L. and *Phaseolus vulgaris* L. (decoration kidney bean) were planted in pots under controlled conditions from suitable soil and irrigation. These plants were left until reaching good vegetative growth (before flowering). We divided plants into four groups: roots, stems, leaves and whole plants. These groups were left in air till complete dryness. Air dried groups of the tested plants were arranged in balls as: roots of *P. major* (100 gm); stems of *P. major* (100 gm); leaves of *P. major* (100 gm); whole plant of *P. major* (100 gm); roots of *P. vulgaris* (100 gm); stems of *P. vulgaris* (100 gm); leaves of *P. vulgaris* (100 gm); whole plant of *P. vulgaris* (100 gm); mixture roots of *P. major* + *P. vulgaris* (50 gm + 50 gm); mixture stems of *P. major* + *P. vulgaris* (50 gm + 50 gm); leaves of *P. major* + *P. vulgaris* (50 gm + 50 gm); mixture whole plant of *P. major* + *P. vulgaris* (50 gm + 50 gm). These balls were hanged (using iron rode) hydroponically in water supplemented with 1, 2, 3, 4 and 5 mM $\text{Pb}(\text{NO}_3)_2$. Samples were tested after 1, 2, 3, 4 and 5 days for their lead ion content under different temperature regimes (10, 20, 30, 40, and 50 °C).

Analysis:

In order to determine the total amount of lead taken up by particular plant organs and whole plants, 0.5-g of the plant material was homogenized in a mortar and pestle with 5- mL of concentrated HNO_3 and concentrated H_2SO_4 (1:1 volume by volume). Then, the extract was warmed until a clear solution formed, 1 ml of H_2O_2 was then added and the extract was heated again. The content of lead in samples was measured with the use of Varian Spectrophotometer AA 20 plus atomic absorption spectrometer equipped with a deuterium lamp for background correction.

Statistical Analysis:

Statistical analyses were carried out using the SPSS BASE 10.0 (SPSS Inc., Chicago, IL) packages, to testify the significant effects of air dry balls of *P. major* and *P. vulgaris* to different concentrations on lead ions. Plant parts uptake ions were subjected to normal procedures of analysis of variance (ANOVA).

RESULTS AND DISCUSSIONS

Results:

In this research, materials were two plants: *P. major* L. and *P. vulgaris* L. (decoration kidney bean). These plants were grown hydroponically in a water medium supplemented with lead ions under different temperature for certain times. The roots of studied plants showed high uptake to large amounts of sublethal 10^{-3} M Pb in water medium (Table 1). Fewer uptakes were observed in the above-ground parts of plants after the third day of treatment with lead ions (Table 2). In comparison to the control *P. major* roots grown under 10°C, the roots of the *P. vulgaris* plants and the mixture roots of both plants changed significantly (Table 3). We observed gradual changes in the Pb-uptake reaching maximum especially, in the 40°C of organs in the plants treated with lead. The absorption of the roots changed gradually and stabilized after the fourth day under the influence of metal ions, from 65% to 79%. The uptake of roots was slowed down in samples treated with other than the 1 mM Pb-ion concentrations and the number of hair roots decreased, which led to lowered water uptake (Table 1). Responses differences between plant organs and whole plants to different Pb-ion values indicate that the lowest concentration of lead ion is the best among the rest of concentrations and has the fastest initiation of the detoxicative system. The *P. major* exhibits the highest sensitivity to lead ions and it has the capacity to take up the largest amount of Pb (66%). We found that roots have the highest accumulation capacity of Pb in the tested plants (Table 1),

Table 1: Removal of lead content (%) by balls of *Plantago major* L. and *Phaseolus vulgaris* L. hanged and left it in water (with 25°C) for five days.

Lead concentrations	Plant parts	1-mM Pb(NO ₃) ₂	2-mM Pb(NO ₃) ₂	3-mM Pb(NO ₃) ₂	4-mM Pb(NO ₃) ₂	5-mM Pb(NO ₃) ₂
Effect of roots ball:						
	<i>Plantago major</i>	77	63	66	69	44
	<i>Phaseolus vulgaris</i>	71	67	61	69	41
	<i>Plantago + Phaseolus</i>	85	76	65	72	55
	LSD (P ≤ 0.05)	5.1	7.8	6.8	4.1	8.6
Effect of stems ball:						
	<i>Plantago major</i>	56	52	51	52	50
	<i>Phaseolus vulgaris</i>	65	54	54	54	45
	<i>Plantago + Phaseolus</i>	55	45	43	41	42
	LSD (P ≤ 0.05)	6.6	9.7	7.9	9.7	6.6
Effect of leaves ball:						
	<i>Plantago major</i>	46	44	42	44	46
	<i>Phaseolus vulgaris</i>	52	40	42	40	42
	<i>Plantago + Phaseolus</i>	50	43	40	43	40
	LSD (P ≤ 0.05)	5.9	7.0	9.9	7.0	5.9
Effect of whole plant ball:						
	<i>Plantago major</i>	66	44	46	44	56
	<i>Phaseolus vulgaris</i>	58	50	50	50	48
	<i>Plantago + Phaseolus</i>	57	50	47	50	47
	LSD (P ≤ 0.05)	9.0	11.2	9.8	11.2	9.0
Effect of plant organs ball:						
	Roots <i>Plantago major</i>	77	63	66	69	44
	Roots <i>Phaseolus vulgaris</i>	71	67	61	69	41
	Stems <i>Plantago major</i>	56	52	51	52	50
	Stems <i>Phaseolus vulgaris</i>	65	54	54	54	45
	Leaves <i>Plantago major</i>	46	44	42	44	46
	Leaves <i>Phaseolus vulgaris</i>	52	40	42	40	42
	LSD (P ≤ 0.05)	8.8	9.7	7.5	9.7	8.8

Table 2: Removal of lead content (%) by balls of *Plantago major* L. and *Phaseolus vulgaris* L. hanged in water (with 25°C) treated with 1 mM of Pb(NO₃)₂.

Treatments duration	Plant parts	After 1-day	After 2-days	After 3-days	After 4-days	After 5-days
Effect of roots ball:						
	<i>Plantago major</i>	68	75	76	77	77
	<i>Phaseolus vulgaris</i>	65	70	71	71	72
	<i>Plantago + Phaseolus</i>	74	75	79	79	79
	LSD (P ≤ 0.05)	6.8	9.8	9.6	6.1	6.1
Effect of stems ball:						
	<i>Plantago major</i>	46	52	55	55	56
	<i>Phaseolus vulgaris</i>	45	54	54	54	54
	<i>Plantago + Phaseolus</i>	44	55	55	55	58
	LSD (P ≤ 0.05)	6.9	6.7	6.9	6.9	6.8
Effect of leaves ball:						
	<i>Plantago major</i>	42	44	55	55	55
	<i>Phaseolus vulgaris</i>	41	42	50	52	55
	<i>Plantago + Phaseolus</i>	40	43	49	50	50
	LSD (P ≤ 0.05)	6.3	7.7	6.9	3.5	3.7
Effect of whole plant ball:						
	<i>Plantago major</i>	55	63	66	66	66
	<i>Phaseolus vulgaris</i>	51	57	60	67	67
	<i>Plantago + Phaseolus</i>	50	60	67	70	70
	LSD (P ≤ 0.05)	9.7	5.4	7.3	6.3	6.3
Effect of plant organs ball:						
	Roots <i>Plantago major</i>	68	75	76	77	77
	Roots <i>Phaseolus vulgaris</i>	65	70	71	71	72
	Stems <i>Plantago major</i>	46	52	55	55	56
	Stems <i>Phaseolus vulgaris</i>	45	54	54	54	54
	Leaves <i>Plantago major</i>	42	44	55	55	55
	Leaves <i>Phaseolus vulgaris</i>	41	42	50	52	55
	LSD (P ≤ 0.05)	4.6	6.4	3.5	5.3	7.2

Table 3: The Removal of lead content (%) by balls of *Plantago major* L. and *Phaseolus vulgaris* L. hanged in water for five days and treated with 1mM of Pb(NO₃)₂.

Water temperaturesPlant parts	10°C	20°C	30°C	40°C	50°C
Effect of roots ball:					
<i>Plantago major</i>	34	62	77	87	53
<i>Phaseolus vulgaris</i>	44	66	71	77	44
<i>Plantago + Phaseolus</i>	49	66	82	91	45
LSD (P ≤ 0.05)	6.4	4.6	5.3	6.2	4.2
Effect of stems ball:					
<i>Plantago major</i>	32	41	54	65	41
<i>Phaseolus vulgaris</i>	31	44	54	64	39
<i>Plantago + Phaseolus</i>	29	43	49	55	42
LSD (P ≤ 0.05)	7.3	6.8	5.6	8.6	9.3
Effect of leaves ball:					
<i>Plantago major</i>	33	39	44	54	38
<i>Phaseolus vulgaris</i>	30	41	46	51	44
<i>Plantago + Phaseolus</i>	36	53	66	68	55
LSD (P ≤ 0.05)	7.1	4.5	6.3	3.6	6.8
Effect of whole plant ball:					
<i>Plantago major</i>	44	49	56	66	55
<i>Phaseolus vulgaris</i>	42	45	56	65	43
<i>Plantago + Phaseolus</i>	51	56	59	62	55
LSD (P ≤ 0.05)	4.4	5.5	6.4	6.4	4.7
Effect of plant organs ball:					
Roots <i>Plantago major</i>	34	62	77	87	53
Roots <i>Phaseolus vulgaris</i>	44	66	71	77	44
Stems <i>Plantago major</i>	32	41	54	65	41
Stems <i>Phaseolus vulgaris</i>	31	44	54	64	39
Leaves <i>Plantago major</i>	33	39	44	54	38
Leaves <i>Phaseolus vulgaris</i>	30	41	46	51	44
LSD (P ≤ 0.05)	5.2	5.0	6.3	4.3	5.4

which is consistent with the earlier data concerning other plants. A steady increase in the level of the biosorped metal was observed in all samples after the first day of treatments until the end of the experimental period when the concentration of lead in roots of *P. major + P. vulgaris* reached 79 % (Table 2). At the same time, roots of *P. major* and *P. vulgaris* uptake lead to a lesser degree (77%, 72%, respectively). After 40°C of water temperature treated with Pb(NO₃)₂, we found about 50% of Pb/ in roots were missing (Table 3).

The presented results showed that about 90-95% of the total amount of lead was localized in roots after the first day of treatments and only 5-10% was transported to it in all after 4-days of lead treatment (Table 2). The level of lead in stems and leaves of all studied plants increased with the time of lead treatment but they were stable in the fourth day (Table 2). In stems, the maximum level of Pb in studied organs is varied. These values were, 58% (*P. major + P. vulgaris*), 56% (*P. major*) and 54% (*P. vulgaris*) and lower than the lead content in the roots by 25%. A significant increase in lead content was observed in leaves of *P. major* and *P. vulgaris* after 3-days compared to the 1-days or 2-days, while it observed in leaves of *P. major + P. vulgaris* after 4-days compared to the 1-days or 2-days (Table 2). The highest amount of lead was found in leaves of *P. major + P. vulgaris* 50% (Table 1). Lower contents of biosorped Pb were found in leaves of bean.

Discussion:

Increasing pollution of the environment caused by heavy metals is becoming a serious problem in the modern world. Plants could be used for phytoextraction, rhizofiltration and/or phytostabilization of heavy metals. Therefore, it is necessary to recognize more broadly plant mechanisms regulating uptake using below or aboveground plant parts, and may lead to reducing the harmful effect of heavy metals.

It is a well-known that roots provide the primary route for the penetration of heavy metals. Roots can take up 3-50 times more lead than stems or leaves (Wozny *et al.*, 1995). The mixture plants we had studied uptake up to 85% Pb in roots and this is similar to results of Tomaszewska *et al.* (1996). Kumar *et al.* (1995) also demonstrated that about 90% of Pb accumulation occurred in roots. The amount of lead ions in the roots of the studied *P. vulgaris* is comparable with the results obtained by Burzynski (1987), who exposed *P. vulgaris*,

cucumber and wheat to 10^{-3} M $PbCl_2$. He showed that cucumber is a plant which accumulates the largest amounts of Pb, but it was very sensitive to this concentration of $PbCl_2$ and, consequently, the plant was killed after only one day of exposition. He also demonstrated that *P. vulgaris* accumulated over 75 mg Pb/g dry wt in roots and exhibited a significant tolerance to this metal (Burzynski, 1987). Antosiewicz and Wierzbicka (1999) localized the highest level of lead in cell walls of *Allium cepa* root tips by means of the conventional electron microscopy preparative technique. Seregin and Ivanov (2001) also obtained similar results in maize treated with 10^{-4} M $Pb(NO_3)_2$. This can be explained by the fact that lead was fixed by functional components of cell walls, such as, among others, polysaccharides (Seregin and Ivanov, 2001). Leopold and Gunther (1997) observed in *Silene vulgaris* cell cultures that only 5% of the total amount of metal ions was present in a soluble extract.

The investigated plants: *P. major* and *P. vulgaris* showed much variation in absorbing lead. The *P. major* biosorped much higher amounts of Pb than *P. vulgaris*. In comparison, the content of this metal in roots of Brassicacea, which is thought to be a moderate uptake of lead, was over 100 mg Pb/g dry wt and about 10 mg Pb/g dry wt in stems, that is; 10 times less than in roots (Kumar et al., 1995). The level of Pb in the roots of plants investigated by Kumar et al. (1995) ranged from 0.82 to 10.9% of dry wt, whereas in our investigations it was 77% of *P. major*, and 71 % of *P. vulgaris*. The best accumulator among the studied samples of the research was the mixture roots of both tested plants recording 85 %.

Biosorption of lead is characteristic to *P. major* than *P. vulgaris*, could have higher phytoextraction ability of lead from polluted water. One can't exclude the fact that genetic manipulations might lead to creating a variety of *P. major*. Moreover, in the other research on lead accumulation in *Brassica juncea*, it has been demonstrated that application of synthetic chelators (Epstein et al., 1999; Blaylock et al., 1997) and hydroponic cultures (Vassil et al., 1998; Salt et al., 1997) raised the concentration of soluble lead and increased its uptake by the plant by creating Pb- complexes. These results suggest that further studies should be conducted on the effect of this chelator on lead uptake by *P. major*. Similarly, a significant increase in lead accumulation occurred in the roots of *Brassica juncea*, however, in the shoots of Indian mustard cultivated in a hydroponic medium (Kumar et al., 1995) it increased significantly only when lead ions concentration reached the value of 100 mg Pb/L in solution.

The gradual uptake of lead in studied organs: roots, stems, and leaves of *P. major* than *P. vulgaris* and reached maximum after the second day and 40°C may suggest that phytochelations play only a transient role in the process of lead ions detoxification. Leopold et al. (1999) observed a similar effect in tomato and *S. vulgaris* cultures exposed to Cd^{2+} and Cu^{2+} ions (0.1 mM), where metal complexes disappeared between the 7th and 14th day of treatment. According to these authors, this proves that the formation of phytochelation-metal complexes plays a part only in the early stage of plant response to stress caused by heavy metal ions, and this is not related to an increased tolerance of plants to metals. Leopold and Gunther (1997) and Leopold et al. (1999) also analyzed extracts of cell cultures and plants and they observed that eluants from HPLC column were complexes of Cd and Cu ions with legends (molecular weight about 13000 Da), which correspond to the phytochelation molecule with oxidised SH-groups. But in tomato and *S. vulgaris* cell cultures exposed to Pb and Zn ions, heavy metals were bound to a lower molecular weight legends (they could not be sufficiently purified for mass spectrometry), but not phytochelations. It is very likely that not only this internal "phytochelatin" detoxification system is connected with the mechanism of plant tolerance to lead, but probably there is also another mechanism, e.g. accumulation on the root surface by binding with polysaccharides (Seregin and Ivanov, 2001) or complexing with organic acids as evidenced by Harmens et al. (1994) in *S. vulgaris* which accumulated zinc.

This study ensure that it is necessary to conduct further research in order to explain the mechanism of high plant tolerance to the effect of lead, as well as to consider the possibility of growing *P. major* varieties with an enhanced ability to uptake Pb, e.g. as a result of adding different accumulative plants which make it possible to increase the uptake of metals with help of different plant parts.

REFERENCES

- Allinson, D.V. and C. Dzialo, 1981. The influence of Pb^{2+} , Cd^{2+} and Ni^{2+} on the growth of rye grass and oats. *Plant Soil*, 62: 81-89.
- Antosiewicz, D. and M.I Wierzbicka, 1999. Localization of lead in *Allium cepa* L. cells by electron microscopy. *J. Microscopy*, 195: 139-146.

Blaylock, M.I., D.E. Salt, S. Dushenkov, O. Zakohrova, C. Gussman, Y. Kapulnik, B.D. Ensley and I. Raskin, 1997. Enhanced accumulation of Pb in Indian mustard by soil-applied chelating agents. *Environ. Sci. Technol.*, 31: 860-865.

Baker, A.I.M., 1995. Metal hyperaccumulation by plants: our present knowledge of the ecophysiological phenomenon. In: 14th Annual Symp. Current Topics in Plant Biochemistry, Physiology and Molecular Biology, Columbia, MO, pp: 7-8.

Bazzaz, F.A., R.W. Carlson and G.L. Rolf, 1974. The effect of heavy metals on plants. I. Inhibition of gas exchange in sunflower by Pb^{2+} , Cd^{2+} , Ni^{2+} and Ti^{2+} . *Environ. Pollut.*, 7: 241-246.

Burzynski, M., 1987. The uptake and transpiration of water and the accumulation of lead by plants growing on lead chloride solutions. *Act. Soc. Bot. Pol.*, 56: 271-280.

Carlson, R.V., F.A. Bazzaz and G.L. Rolf, 1995. The effect of heavy metals on plants. II. Net photosynthesis and transpiration of hole corn and sunflower plants treated with Pb^{2+} , Cd^{2+} , Ni^{2+} and Ti^{2+} . *Environ. Res.*, 10: 113-120.

Epstein, A.L., C.D. Gussman, M.I. Blaylock, U. Yerrniyahu, I.W. Huang, Y. Kapulnik and C.S. Orser, 1999. EDTA and Pb- EDTA accumulation in *Brassica juncea* grown in Pb-amended soil. *Plant & Soil*, 208: 87-94.

Grill, E., S. Lomer, E.L. Winnaeker and M.H. Zenk, 1989. Phytochelatins, the heavy-metal binding peptides of plants, are synthesized from glutathione by a specific γ -glutamyl dipeptidyl transpeptidase (phytochelatin synthase). *Proc. Natl. Acad. Sci. USA*, 86: 6838-6842.

Kucharski, R., A. Sas-Nowosielska, M. Pogrzeba, K. Krynski and E. Malkowski, 1999. Perspectives of using phytoextraction for soil cleaning in Polish conditions. *Protection of Environment*, 18: 469-475.

Kumar, P.B.A.N., V. Dushenkov, H. Motto and I. Raskin, 1995. Phytoextraction: the use of plants to remove heavy metals from soils. *Environ. Sci. Technol.*, 29: 1232-1238.

Leopold, I. and D. Gunther, 1997. Investigation of binding properties of heavy-metal-peptide complexes in plant cell cultures using HPLCICP-MS. *J. Anal. Chem.*, 359: 364-370.

Leopold, I., D. Gunther, I. Schmit and D. Neumann, 1999. Phytochelatins and heavy metal tolerance. *Phytochemistry*, 50: 1323-1328.

Maitani, T., H. Kubota, K. Sato and T. Yamada, 1996. The composition of metals bound to class III metallothionein (phytochelatin and its desglycyl peptide) induced by various metals in root cultures of *Rubia tinctorum*. *Plant Physiol.*, 110: 1145-1150.

Morel, I.L., G. Bitton, C. Schwartz and M. Schiavon, 1997. Bioremediation of soils and waters contaminated with micropollutants: Which role for plants? In: Zelikoff, J.T. (Ed.), *Ecotoxicology: Responses, Biomarkers and Risk Assessment*. SOS Publications, Fair Haven, New York, USA, pp: 37-74.

Salt, D.E., I.J. Pickering, R.C. Prince, D. Gleba, S. Dushenkov, R.D. Smith and I. Raskin, 1997. Metal accumulation in aquacultured seedlings of Indian mustard. *Environ. Sci. Technol.*, 31: 1631-1644.

Seregin, T.V. and V.B. Ivanov, 2001. Physiological aspects of toxic action of cadmium and lead on high plants. (in Russian). *Fiziol. Rasl. (Plant Physiology) (Moscow)*, 48: 606-630.

Sheoran, I.S. and R. Singh, 1993. Effect of heavy metals on photosynthesis in higher plants. In: Abml, Y.P., Mohanty P. and Govindjee, A. (Eds.), *Photosynthesis: Photoreactions to Plant Productivity*, Oxford and IBH Publishing Co. PVI. Ltd, New Delhi, Bombay., pp: 453-468.

Stiborova, M., M. Ditrichova and A. Brezinova, 1987. Effect of heavy metal ions on growth and biochemical characteristics of photosynthesis of barley and maize seedlings. *Biol. Plant.*, 29: 453-467.

Stiborova, M., M. Doubravova, A. Brezinova and A. Friedrich, 1986. Effect of heavy metal ions on growth and biochemical characteristics of photosynthesis of barley (*Hordeum vulgare* L.). *Photosynthetica*, 20: 418-425.

Titov, A.F., V.V. Talanova and N.P. Boeva, 1996. Growth regulation of barley and wheat seedlings to lead and cadmium. *Biol. Plant.*, 38: 431-436.

Tomaszewska, B., A. Tukendorf and D. Baralkiewicz, 1996. The synthesis of phytochelatins in lupin roots treated with lead ions. *The Science of Legumes*, 3: 206-217.

Vassil, A.D., Y. Kapulnik, I. Raskin and D.E. Salt, 1998. The role of EDTA in Pb transport and accumulation by Indian mustard. *Plant Physiol.*, 117: 447-453.

Wozny, A., M. Krzeslowska and B. Tomaszewska, 1995. Heavy metals. In: Wozny, A. (Ed.), *Lead in Plant Cells*. Sorus, Poznan, Poland, pp: 21-110.